Estonia's growing season has lengthened by three to four weeks since 1965, mostly in spring

By Hanna Vesterinen, Chronicle Science Desk · 14 min read

Of those botanists' 1,600 study of Estonian spring leaves, 12,140 trees nearby, and 1,200 beehives, the most striking finding is how much earlier spring arrives. The average first day of leaf-out on the 1,600 monitored trees has moved forward by an average of 21 days, or roughly three weeks, between the 1965–1974 baseline and the 2015–2024 decade. Some species advanced by another week on top of that. Honey harvests, which depend on a long stretch of warm weather after the trees flower, have grown by about 40 percent over the same period. The data are unusually clean because the same observers, using the same methods, have walked the same routes for more than half a century. Few countries can match that record. Estonia can, and the result is a rare piece of public science that tells ordinary people what is changing in the woods around their own villages.

Published 14 March 2026 · Updated 15 March 2026

At a glance Spring arrives on average three weeks earlier than it did in the 1960s, and the season is now about a month longer overall. Honey harvests have grown by roughly 40 percent. The trend is driven mostly by warmer March and April temperatures, and it is consistent across nearly every species the network tracks.

The season really has grown by about a month

The simplest way to see the change is to compare the start of spring with the end of it. Spring, here, is the stretch of days when the average temperature sits between 5 °C and 15 °C. In the 1965–1974 baseline, that stretch began on April 18 and ended on May 23, a little over five weeks. In the most recent decade, it began on March 28 and ended on June 4, more than nine weeks. The difference is not a rounding error. It is the kind of shift that changes when you can plant potatoes, when you can safely leave a greenhouse unheated, and when a beekeeper has to be ready to move hives.



Leaf-out has moved forward by an average of 4.2 days per decade, and flowering by 3.5 days per decade. Autumn leaf-fall is the only phenology that has not shifted, and it is the one most likely to change next.

The trend is not smooth. Some years are wildly out of step with the decade around them, and a few of the most striking ones are worth remembering. 1989 opened with a March that was 4.1 °C above the baseline mean, and the first leaves on the monitored birches appeared 19 days earlier than the previous record. 2003 was the opposite: a cold April pushed leaf-out back by nearly two weeks, and the honey harvest that year was the smallest in the network's history. 2018 and 2020 both produced warm springs but for different reasons; 2018 was dry, 2020 was wet, and the trees responded to the temperature rather than the rainfall.

Why the trees are waking up

The driver is straightforward. Estonian March and April temperatures have risen by about 1.9 °C since the baseline, and the trees are tracking that warmth almost one-for-one. Regression of leaf-out date against spring temperature gives a slope of roughly −4.6 days per °C, which is what you would expect if nothing else were changing. Something else is changing, though. Winter chilling has also declined, and some species may be starting to need more heat to wake up than they used to.

> "We are not just counting dates. We are watching the trees answer a question we asked them on purpose, and the answer is getting clearer every year."
>
> — Dr. Kristiina Mägi, lead author, Estonian Phenology Network

The honey numbers are the most economically vivid. Beekeepers in the network recorded an average harvest of 38 kg per hive in the 1965–1974 decade and 53 kg per hive in the most recent decade, a rise of about 40 percent. The increase is not because bees have become more productive. It is because the nectar flow now arrives earlier, lasts longer, and often overlaps with warmer, calmer weather that lets foragers work more hours. Beekeepers who moved hives into the network's monitored apiaries in the 1970s and 1980s have watched the change in real time.

Which species are moving fastest

Not every tree is responding at the same speed. Some species are far more sensitive to spring warmth than others, and the differences matter for anyone trying to predict what the forest will look like in 2040.

Species	Change in leaf-out (days per decade)	Change in flowering (days per decade)
Silver birch (*Betula pendula*)	−5.1	−4.4
Pedunculate oak (*Quercus robur*)	−3.8	−3.0
European aspen (*Populus tremula*)	−6.2	−5.5
Rowan (*Sorbus aucuparia*)	−4.0	−3.6
Bird cherry (*Prunus padus*)	−4.7	−4.1
Hazel (*Corylus avellana*)	−2.9	−2.4
Norway spruce (*Picea abies*)	−1.6	−1.2
Scots pine (*Pinus sylvestris*)	−1.1	−0.8

Aspen and bird cherry are the fastest responders in the network, while the conifers are the slowest. That gap is widening the mismatch between when insects emerge and when the flowers they pollinate are open. Pollinators that have not yet adjusted to the earlier season are showing up to a meal that is already half over.

Not a trend in a century: nearly four weeks in 61 years on the region's lighter, sandier soils

The 1965–2024 record is unusual for a reason that matters for how far it can be stretched. Most long phenology records in Europe begin in the 1990s, after the warming had already accelerated. The Estonian network started in 1964 and ran through the cold decades, so it captures the full shape of the change rather than only its steep end. That is why the trend can be read as a century-scale signal rather than a short-term fluctuation.



Mean leaf-out day across the 1,600 monitored trees, 1965–2024. The dashed line is the fitted trend: 19.7 days earlier per decade, or roughly 3.9 weeks over the full record.

The strength of the trend varies by soil. Trees on the network's lighter, sandier plots have moved by 4.8 days per decade, while those on heavier, clay-rich soils have moved by 2.6 days per decade. The difference is consistent with soils that warm and dry faster in spring letting roots access water earlier in the season. It is a small but telling detail that suggests the trees are responding to more than just the air temperature above them.

Where the change is strongest

The network's 312 plots are spread across the country, from the dunes of Pärnu to the inland forests of Võrumaa. The earliest leaf-out dates are now recorded in the southwest, where the maritime climate has always been mildest, and the latest in the northeast, near the Russian border. The gradient between those two corners has not disappeared; it has simply moved forward in time.

Region	Mean leaf-out, 1965–1974	Mean leaf-out, 2015–2024	Change
Southwest (Pärnu, Lääne)	21 May	1 May	−20 days
Central (Tartu, Viljandi)	24 May	4 May	−20 days
Northeast (Ida-Viru, Lake Peipsi)	28 May	9 May	−19 days
Southeast (Võrumaa)	26 May	6 May	−20 days
Islands (Saaremaa, Hiiumaa)	19 May	28 April	−21 days

Mean leaf-out day for silver birch, by region. The islands warm earliest and have moved the most in absolute terms.

Honey, pollen, and the wider ecosystem

The honey harvest is the most visible consequence, but it is not the only one. Pollen counts in the network's air samplers have begun rising earlier and falling later, which means the allergy season is now about 11 days longer than it was in the 1970s. Hazel pollen, which is the first major allergen of the year, has shifted the most. Frost damage to flowers is also changing. Late spring frosts that once killed a large share of the bird cherry crop have become rarer, though they have not disappeared.

Year	Honey harvest (kg per hive)	Notable conditions
1965	36	Cool, wet spring; poor nectar flow
1974	41	Average season
1989	49	Warm March; first 19-day-early leaf-out
2003	31	Cold April; smallest harvest in the network
2010	47	Long, dry summer
2018	52	Dry spring; trees responded to heat, not rain
2020	58	Wet spring; record honey year
2024	53	Second-warmest year in the network

Honey harvest per hive, 1965–2024. The 2020 figure is the highest on record; 2003 is the lowest.

What the network still cannot say

The data are clear about what has happened, but they are less clear about why some species are responding more than others. The network records dates, not mechanisms. Researchers can infer that aspen is more sensitive to warmth than spruce, but they cannot yet say whether that is because of differences in bud structure, in chilling requirements, or in something else entirely. Those questions require experiments, not observation, and the network is now being asked to support them.

There is also a limit to how far the trend can be extrapolated. The relationship between spring warmth and leaf-out is roughly linear over the range of temperatures the network has seen, but it will not stay linear forever. Trees cannot leaf out before the end of winter, and at some point the chilling they need to wake up will no longer be reliably delivered. The network's data are not yet close to that limit, but they are close enough that it is worth watching for.

Why the network matters now

Phenology is one of the few fields in which ordinary people can contribute data that professionals actually use. The Estonian network has always relied on volunteers, and its observers are now among the most experienced in the country. Their records are the reason the trend can be read with confidence rather than guesswork. As the climate keeps changing, the value of that long, boring, careful record will only grow.

Sources

1. Mägi, K., Vesterinen, H., & Tamm, R. (2025). Three weeks earlier: 60 years of spring leaf-out in Estonian forests. *Estonian Phenology Network Annual Report*, 41, 1–88.
2. Vesterinen, H. (2026, March 14). Estonia's growing season has lengthened by three to four weeks since 1965, mostly in spring. *The Chronicle Science*.
3. Kask, L., & Pärnoja, E. (2024). Honey harvests and spring weather in the Estonian beekeeping record, 1965–2024. *Journal of Apicultural Science*, 68(2), 211–230.
4. Ilves, A., Rannik, Ü., & Sepp, M. (2023). Pollen seasons are getting longer: 30 years of aerobiological monitoring in Tartu. *Aerobiologia*, 39(4), 655–671.
5. Tõnisson, J., & Kuusk, E. (2022). Frost damage to bird cherry blossom in Estonia, 1965–2020. *Forestry Studies*, 76, 5–24.
6. Mägi, K., & Vesterinen, H. (2025). Soil texture and the timing of spring in Estonian forests. *Estonian Phenology Network Annual Report*, 41, 89–112.
7. Vesterinen, H. (2026, March 15). Correction: honey harvest figures, 1989 and 2003. *The Chronicle Science*.

How this was checked

The author is Hanna Vesterinen, Chronicle Science Desk. Reporting was supported by a grant from the Estonian Science Foundation (grant PUT-2241). The author has no financial relationship with the Estonian Phenology Network.

Data and code for the figures in this article are available at github.com/chronicle-science/estonia-leaf-out-2026. Raw observer records are held by the Estonian Phenology Network and can be requested for research use.

Corrections and changes

15 March 2026 — Corrected honey harvest figures for 1989 and 2003. The first version reported 44 kg and 28 kg per hive; the correct values are 49 kg and 31 kg per hive.

14 March 2026 — Original publication.